

**TuBAvi (2017-20)**  
**TuBAvi-2 (2021-24)**

Collective projects within the poultry sector funded with the support of the **European Agricultural Fund for Rural Development (EAFRD)**

[https://ec.europa.eu/agriculture/rural-development-2014-2020\\_en](https://ec.europa.eu/agriculture/rural-development-2014-2020_en)

**Ministry of agriculture, food sovereignty and forestry** –  
National Rural Development Programme 2014/2022 – Measure 10.2 –  
Conservation, use and sustainable development of genetic resources  
in agriculture



**Project coordinator**  
Prof. Silvia Cerolini  
Department of Veterinary Medicine and Animal Sciences  
University of Milan  
Email [silvia.cerolini@unimi.it](mailto:silvia.cerolini@unimi.it)  
[www.pollitaliani.it](http://www.pollitaliani.it)

**CONSERVATION OF BIODIVERSITY IN ITALIAN POULTRY BREEDS:**  
deepening and monitoring  
**TuBAvi-2**



**Breed data sheet**

**ROBUSTA LIONATA**  
*Gallus gallus domesticus Sp.*

**Origin and morphological,  
genetic, reproductive,  
and productive traits**



**FONDO EUROPEO AGRICOLO PER LO SVILUPPO  
RURALE: l'Europa investe nelle zone rurali**



**MINISTERO DELL'AGRICOLTURA  
DELLA SOVRANITÀ ALIMENTARE  
E DELLE FORESTE**





The presented data were registered in nucleus populations of Robusta Lionata conserved at the “Sasse Rami” Experimental Farm, in Ceregnano (Rovigo).

Latest update: October 14<sup>th</sup>, 2023



## Robusta lionata

*Gallus gallus domesticus Sp.*

Breed data sheet: origin and morphological, genetic, reproductive, and productive traits

### Breed origin and development

Name of the breed	Robusta lionata
Synonyms or local names	-
Geographic origin	Veneto (Rovigo)
Geographic distribution	Veneto
Estimated total population size	452 (Castillo et al., 2021)
Extinction risk status (FAO, 1998)	Threatened conserved
Any other specific information	-

<b>Historical origin</b>
The Robusta Lionata breed was created in 1965 at the Poultry Experimental Station of Rovigo. It was created using the Tawny Orpington and White America breeds during the selection process. The breed is characterised by good egg and meat production, and it is reared for niche products. Hens show high aptitude for broodiness and parental care, also towards eggs and chicks of other species. It is included as a local breed in the National Plan on Biodiversity in Agriculture and in the Atlas of Traditional Agri-food Products ( <i>Atlante dei Prodotti Agroalimentari Tradizionali</i> ) of the Veneto region.

## Qualitative and quantitative morphological traits in adult breeders

### Discrete or qualitative traits

Feather morphology	Normal
Feather distribution	Normal
Plumage structure	Abundant, slightly soft
Plumage colour	Tawny
Colour features	Bi-colour, with sexual dimorphism
Chick plumage colour	Tawny down with little brown spots on the head
Comb type	<b>Simple comb</b> , upright
Comb spikes	Five or six spikes
Ear-lobe colour	Red
Beak colour	Orange
Iris colour	Orange to red
Muffs	Absent
Beard	Absent
Tuft	Absent
Skin colour	Yellow
Shank colour	Yellow
Shank feathering	Free from feathers
Skeletal variants	-
Other specific and distinct visible traits	-

<b>Colour pattern</b>
In the <b>male</b> , yellow-tawny of a warm tone, hackle feathers with black and irregular spots or pencilling, flight feathers with brown veining. In the <b>female</b> , slight patterning on the back is admitted. All the tail feathers black at the end, with green luster.

### Quantitative traits

Parameters	Male		Female	
	Average	Min-max	Average	Min-max
Body weight (g)	3950	3620-4280	2750	2120-3010
Body length (cm)	47	45-49	41	38-45
Chest circumference (cm)	41	37-50	37	33-43
Shank length (cm)	11	9-14	9	8-11
Shank diameter (cm)	6	5-6	5	4-5
Wing span (cm)	53	51-56	47	43-49

### Robusta lionata male and females



UniPD



UniPD

### Genetic traits

#### Characterisation of the breed with Single Nucleotide Polymorphisms (SNPs)

Molecular marker	Affymetrix Axiom 600K Chicken Genotyping Array
Laboratory that performed the analyses	Department of Agronomy, Food, Natural Resources, Animals and Environment (DAFNAE) University of Padua
Analysed parameters	MAF: minor allelic frequency Ho: observed heterozygosis He: expected heterozygosis F <sub>HOM</sub> : inbreeding coefficient

Year		N**	MAF	Ho	He	F <sub>HOM</sub>
2019	Mean	23	0.305	0.181	0.185	0.508
	SD*		0.345	0.199	0.195	0.039

\*SD: standard deviation; \*\*N: number of samples

#### Characterisation of nucleus populations with microsatellites

Molecular marker	Microsatellites (26 markers)
Laboratory that performed the analyses	Laboratory of Animal Molecular Genetics Department of Veterinary Science (DSV) University of Turin
Analysed parameters	Ne: effective number of alleles Na: observed number of alleles I: Shannon diversity index H-Ind: individual variability index Ho: observed heterozygosis (average H-Ind) He: expected heterozygosis F: fixation index P: average kinship index

Year		N**	Na	Ne	I	Ho	He	F	P
2020	Mean	23	2.857	1.838	0.689	0.345	0.393	0.101	0.66
	SE*		0.294	0.161	0.104	0.054	0.058	0.059	

\*SE: standard error; \*\*N: number of samples

## Reproductive and productive quantitative traits

### Oviposition, brooding and incubation data

Age at sexual maturity of hens (weeks)	22-27
Length of first oviposition cycle (weeks)	N.a.**
Annual egg production per hen (min-max)*	160-170
Average clutch size (min-max)	N.a.**
Clutch interval (days)	N.a.**
Incubation length (days)	21

\*As measured during the first year of age, min-max of family line

\*\*N.a.: Not available information

### Egg-quality traits

Parameters	First oviposition cycle	
	Average	Min-max
Egg weight (g)	57,5	Not available
Shell colour	Pinkish	

Parameters (sample measurement)	Average	Min-max
Egg weight (g)	60.6	57.2-64.0
Shell weight (g)	5.41	4.94-5.88
Albumen weight (g)	36.6	33.1-40.1
Yolk weight (g)	17.4	16.1-18.7
Egg Shape Index*	0.76	0.73-0.80

\* Egg Shape Index (ESI) = short diameter/long diameter x 100

### Reproductive traits

Incubation parameters	First oviposition cycle	
	Average	Min-max*
Fertility (% produced eggs)	81	74-85
Hatchability (% fertile eggs)	74	68-76
Hatchability (% produced eggs)	60	57-63

\*Per family line

### Slaughter data (age: 24 weeks)

Slaughter parameters	Average	
	Male	Female
Live weight (g)	3393	2407
Carcass weight (eviscerated) (g)	2219	1550
Carcass weight (eviscerated) yeald (%)	65.4	64.4

## Rearing traits

Breed type	Rustic
Growth speed (precocious vs tardive)	Tardive
Feathering speed (precocious vs tardive)	Precocious
Broodiness	Yes
Parental care attitude	Yes
Ease of breeding	Yes
Male:female ratio for breeding	1:10
Tolerance or resistance to diseases and parasites	Yes
Tolerance to extremes of temperature	Yes
Reported uses (meat, eggs)	Primary: meat Secondary: eggs